

Ophthalmic Physicians and Surgeons

2nd Floor, 87 Wickham Terrace BRISBANE QLD 4000

Phone: 07 3831 0101 Fax: 07 3831 3203

www.terraceeyecentre.com.au Email: admin@terraceeyecentre.com.au

# Dr Sunil Warrier

## Cataract Surgery - Post - operative Instructions

#### **IMPORTANT**

- Avoid heavy rubbing or any bumping of the eye for the first month after surgery.
- Avoid sustained bending or lifting of heavy objects.
- You should not experience any pain beyond some mild irritation of discomfort.
- If you do feel pain contact Dr Warrier's surgery immediately.
- Panadol or Panamax should be all that is necessary to ease your discomfort.

#### **EYE DROPS**

 You have three bottles of eye drops to use, CHLORSIG, ACULAR and PREDNEFRIN FORTE (this one must be shaken well prior to use)

#### The first four weeks: / /

- Instil ONE drop from each bottle into the operated eye 4 times a day.
- Easy times to remember are at breakfast, lunch, dinner and prior to going to bed.
- The order of the drops is not important.
- After instilling the drop, gently close your eye for 30 seconds and wait at least 2 minutes before using the the next drop (remembering to shake it first).

#### **HYGIENE**

- Without putting excessive pressure on the eyeballs, clean eyelashes daily by rubbing gently with fingertips while showering.
- Avoid soap or shampoo in the eye
- No eye makeup for 1 week
- Wear the eye shield at night for the first 1 week.

### **ACTIVITIES**

- DO NOT DRIVE until instructed by Doctor.
- No Swimming for 1 month.
- You can return to your normal activities (eg bowls) without concern.

#### GLASSES

- You may use your current glasses until you visit your Optometrist in four weeks.
- If you normally wear dark glasses, continue to do so if you wish.

### **FOLLOW UP**

• If your eye becomes more red, more uncomfortable, painful of the vision becomes worse, please contact Dr Warrier for a check immediately.

Business Hours: 07 3831 0101